Chapter-3.

NEUROPHYSIOLOGY OF SACCADES, PURSUITS, AND VERGENCE

Dr Partha Chowdhury, Dr Brinda Shah

ABSTRACT

The human visual system relies on coordinated ocular motor mechanisms to maintain stable, accurate, and efficient vision. Among these mechanisms, saccades, smooth pursuit, and vergence eye movements represent the three principal modes of ocular motor control, enabling rapid shifts in gaze, continuous tracking of moving targets, and binocular alignment for depth perception, respectively. These eye movements are regulated by highly specialized and interconnected neural networks involving the cerebral cortex, cerebellum, brainstem, and subcortical structures. Understanding the neurophysiological basis of these ocular motor systems is essential for interpreting normal visual function as well as diagnosing neurological and neuro-ophthalmic disorders. This chapter provides a comprehensive review of the neurophysiological mechanisms underlying saccadic, smooth pursuit, and vergence eye movements, emphasizing their distinct functional characteristics, shared neural pathways, and regulatory mechanisms. Furthermore, it discusses their clinical relevance, associated neurological disorders, and implications for neuro-optometric assessment and rehabilitation, highlighting their significance in both clinical practice and vision science.

Keywords: Saccadic eye movements; Smooth pursuit; Vergence eye movements; Ocular motor control.

3.1 INTRODUCTION

Precise control of eye movements is fundamental to maintaining clear, stable, and single binocular vision. The human visual system is highly dynamic, requiring the eyes to continuously adjust their position to accurately detect, track, and fixate on objects within a constantly changing environment. These adjustments are achieved through three principal types of eye movements: saccades, which rapidly redirect the line of sight between fixation points; smooth pursuit movements, which enable continuous tracking of moving objects; and vergence movements, which coordinate the alignment of both eyes when viewing objects at different distances (Leigh & Zee, 2015; Carpenter, 1988). Together, these ocular motor systems ensure optimal visual acuity, depth perception, and spatial orientation during everyday activities such as reading, driving, and object manipulation. The neurophysiology underlying these eye movements involves a sophisticated network of cortical, subcortical, brainstem, and cerebellar structures that integrate visual, vestibular, proprioceptive, and cognitive information to produce precise motor commands. Regions such as the frontal and parietal eye fields, superior colliculus, brainstem gaze centres, basal ganglia, and cerebellum work in close coordination to initiate, execute, and continuously refine ocular movements (Kandel et al., 2021). This integrated control allows the visual system to maintain stable fixation, accurately follow moving targets, and preserve binocular alignment despite constant head and body movements. Understanding the neurophysiology of saccades, smooth pursuits, and vergence is essential not only for appreciating normal visual function but also for recognizing the mechanisms underlying numerous neuro-ophthalmic and neurological disorders. Dysfunction of these ocular motor pathways may result in abnormalities such as gaze palsies, impaired smooth pursuit, convergence insufficiency, nystagmus, and other eye

movement disorders that significantly affect visual performance and quality of life. Advances in neuroimaging, electrophysiology, eye-tracking technology, and neurorehabilitation have considerably enhanced our understanding of these complex neural systems, facilitating improved diagnosis, targeted therapy, and patient care (Leigh & Zee, 2015).

3.2 NEUROPHYSIOLOGY OF SACCADES

Saccades are among the most extensively studied eye movements in neuroscience due to their remarkable speed, precision, and dependence on complex neural circuits. They are rapid, conjugate, ballistic eye movements that redirect the line of sight so that the image of an object of interest falls on the fovea, the retinal region responsible for maximum visual acuity. Saccades occur voluntarily during visual exploration and reading, as well as reflexively in response to sudden peripheral stimuli. The neurophysiology of saccades involves coordinated activity within cortical, subcortical, cerebellar, and brainstem structures that integrate sensory information, motor planning, and execution to generate highly accurate eye movements.

A. Characteristics of Saccades: Saccades are rapid, ballistic eye movements that shift the eyes from one point of fixation to another, enabling efficient visual exploration of the environment. Unlike smooth pursuit movements, saccades are preprogrammed before initiation, meaning their direction and velocity cannot be altered once the movement begins. Humans typically generate 2–5 saccades per second, allowing continuous scanning of the surroundings, reading, and rapid acquisition of new visual information. The speed of a saccade increases with its amplitude, with peak velocities ranging from 100°/s for small saccades to over 700°/s for larger movements, making them among the fastest voluntary movements of the human body (Carpenter, 1988; Leigh & Zee, 2015).

During saccadic eye movements, visual sensitivity is briefly reduced through a process known as saccadic suppression, which prevents the perception of motion blur and ensures stable vision. Saccades play an essential role in redirecting gaze toward objects of interest, exploring complex visual scenes, reading, shifting visual attention, acquiring peripheral targets, and maintaining binocular alignment. Based on their mode of initiation, saccades are classified as reflexive, voluntary, memory-guided, predictive, and express saccades, each involving specialized neural pathways responsible for visual processing, motor planning, and cognitive control.

B. Cortical and Subcortical Control of Saccades: The generation of saccadic eye movements requires precise coordination among multiple cortical and subcortical structures. Rather than originating from a single anatomical center, saccades emerge through interactions between sensory, cognitive, and motor networks that collectively determine when, where, and how rapidly the eyes should move.

Frontal Eye Fields (FEF): The frontal eye fields, located within the posterior portion of the frontal cortex (Brodmann area 8), are the primary cortical centers responsible for voluntary saccade initiation. Neurons within the FEF encode movement direction, amplitude, and target location before movement begins. Electrical stimulation of the frontal eye fields produces contralateral saccades, whereas lesions result in impaired voluntary gaze shifts, prolonged reaction times, and difficulty initiating intentional eye movements. Beyond motor execution, the FEF also contributes to selective visual attention and predictive eye movement planning, allowing individuals to anticipate target motion before movement occurs (Leigh & Zee, 2015).

Supplementary Eye Fields (SEF): Located on the medial frontal cortex, the supplementary eye fields coordinate internally generated and sequential eye movements. Unlike the frontal eye fields, which primarily respond to external visual stimuli, the SEF contributes to movement planning, motor sequencing, learning, and memory-guided saccades. The supplementary eye fields are particularly important during complex visual tasks requiring multiple sequential gaze shifts, such as reading, visual search, and navigation through unfamiliar environments.

Posterior Parietal Cortex (PPC): The posterior parietal cortex integrates visual, proprioceptive, and spatial information to determine the location of potential targets within the external environment. It plays a central role in visuospatial attention by selecting the most behaviorally relevant stimulus among multiple competing visual inputs. Damage to the posterior parietal cortex may result in impaired target selection, delayed saccade initiation, and hemispatial neglect, in which patients fail to attend to objects located within one side of visual space.

Superior Colliculus: The superior colliculus, situated within the dorsal midbrain, represents one of the most important subcortical centers controlling saccadic eye movements. It integrates visual, auditory, and somatosensory information into motor commands that guide eye movements toward behaviorally significant targets. The superior colliculus contains a topographically organized map of visual space, allowing precise encoding of movement direction and amplitude. It receives input

from the retina, visual cortex, frontal eye fields, and basal ganglia while projecting to brainstem premotor nuclei responsible for executing saccades. Besides initiating reflexive saccades, the superior colliculus also participates in visual attention, multisensory integration, and orienting behaviors (Sparks, 2002).

C. Brainstem Mechanisms of Saccadic Eye Movements: The final motor commands responsible for saccadic eye movements are generated within specialized brainstem premotor centers that activate extraocular muscles through cranial nerve nuclei.

Horizontal Saccades: Horizontal saccades are controlled primarily by the paramedian pontine reticular formation (PPRF) located within the pons. The PPRF contains excitatory burst neurons that generate high-frequency neural discharges immediately before horizontal eye movements. These neurons activate the ipsilateral abducens nucleus, whose motor neurons innervate the lateral rectus muscle. Simultaneously, internuclear neurons within the abducens nucleus project through the medial longitudinal fasciculus (MLF) to the contralateral oculomotor nucleus, activating the medial rectus muscle and producing coordinated conjugate gaze.

Vertical Saccades: Vertical saccades originate from the rostral interstitial nucleus of the medial longitudinal fasciculus (riMLF) located within the upper midbrain. Neurons within the riMLF project bilaterally to the oculomotor and trochlear nuclei, coordinating elevation and depression of both eyes. Adjacent structures, including the interstitial nucleus of Cajal, contribute to vertical gaze holding and stabilization following movement completion.

Burst Neurons and Omnipause Neurons: Precise control of saccadic timing depends upon interactions between burst neurons and omnipause neurons.

- Excitatory burst neurons generate the high-frequency motor commands required for rapid eye acceleration.
- Inhibitory burst neurons suppress antagonist extraocular muscles, ensuring smooth execution.
- Omnipause neurons, located within the pontine raphe nuclei, remain continuously active during fixation, preventing unwanted saccades. Immediately before saccade initiation, omnipause neurons cease firing, allowing burst neurons to activate ocular motor nuclei. Once the movement is completed, omnipause neuron activity resumes, terminating the saccade and restoring stable fixation.

This highly coordinated neuronal interaction ensures accurate initiation, execution, and termination of every saccadic eye movement.

D. Cerebellar Modulation of Saccades: Although cortical and brainstem structures generate the primary motor commands for saccades, the cerebellum ensures their accuracy, timing, and coordination through continuous error correction and motor adaptation. The oculomotor vermis regulates saccadic amplitude and timing, while the fastigial nucleus controls movement acceleration, deceleration, and endpoint precision. By comparing intended and actual eye movements, the cerebellum continuously adjusts future saccades to maintain accurate gaze despite fatigue, aging, or neurological changes. Damage to cerebellar pathways results in saccadic dysmetria, causing hypometric (undershooting) or hypermetric (overshooting) saccades and impaired visual stability (Robinson, 1981).

.

3.3. NEUROPHYSIOLOGY OF SMOOTH PURSUIT

A. Characteristics of Smooth Pursuit Movements: Smooth pursuit eye movements are specialized, slow, conjugate eye movements that enable continuous tracking of moving objects while maintaining their image on the fovea, the region of the retina responsible for the highest visual acuity. Unlike saccades, which rapidly redirect the line of sight between stationary targets, smooth pursuit movements continuously match the velocity and direction of a moving object, thereby minimizing retinal image slip and preserving clear vision during motion. These movements are essential for many everyday visual tasks, including driving, reading moving text, playing sports, observing moving vehicles, and following people or animals in motion. Smooth pursuit movements typically occur at velocities below 100°/s, although they may occasionally reach higher velocities depending on the speed of the moving target. When target velocity exceeds the tracking capability of the pursuit system, corrective "catch-up" saccades are generated to reposition the fovea accurately on the target (Carpenter, 1988; Leigh & Zee, 2015).

Unlike saccadic eye movements, smooth pursuit is entirely dependent on the presence of a moving visual stimulus and cannot normally be initiated voluntarily in the absence of retinal image motion. This dependence on continuous visual feedback distinguishes pursuit movements from other ocular motor systems and highlights their reliance on specialized neural mechanisms dedicated to motion detection and processing (Krauzlis, 2004). The effectiveness of smooth pursuit is commonly assessed by measuring pursuit gain, defined as the ratio of eye velocity to target velocity. An ideal

pursuit system exhibits a gain close to one, indicating accurate matching between eye and target movements, whereas reduced gain results in retinal slip and frequent corrective saccades.

B. Cortical Mechanisms: The initiation and maintenance of smooth pursuit eye movements depend on a distributed network of cortical regions that process visual motion and transform sensory information into coordinated motor commands. The primary cortical center responsible for motion perception is the middle temporal visual area (MT or V5), which contains highly specialized neurons that detect the direction and speed of moving objects. Visual information originating from the retina is transmitted through the lateral geniculate nucleus to the primary visual cortex before reaching area MT, where motion-sensitive neurons analyze the velocity, direction, and characteristics of target movement. Adjacent to MT is the medial superior temporal area (MST), which further processes complex motion patterns, including optic flow generated during self-motion and large-field visual movement. The MST integrates information regarding target trajectory, acceleration, and anticipated motion, enabling predictive tracking even when the moving object is temporarily obscured from view. Together, MT and MST provide continuous sensory information that forms the basis for smooth pursuit initiation and maintenance (Ilg, 2008; Thier & Ilg, 2005). In addition to these visual motion areas, the frontal eye fields (FEF) play a significant role in voluntary pursuit initiation, predictive tracking, and adjustment of pursuit velocity. The FEF contributes particularly when target movement is predictable, allowing the ocular motor system to initiate pursuit before visual feedback alone would be sufficient. This predictive capability minimizes retinal image slip and enhances tracking accuracy during repetitive or anticipated target motion (Leigh & Zee, 2015). Collectively, these cortical regions integrate visual perception, attention, and motor planning to generate smooth and coordinated pursuit eye movements.

C. Brainstem and Cerebellar Circuits: The smooth pursuit pathway involves close coordination between the brainstem and cerebellum to ensure accurate and stable eye tracking. Visual motion signals from the middle temporal (MT) and medial superior temporal (MST) areas are transmitted through the dorsolateral pontine nuclei (DLPN) and the nucleus reticularis tegmenti pontis (NRTP) to the flocculus and paraflocculus of the cerebellum. These cerebellar regions regulate pursuit gain, movement accuracy, and adaptive control by continuously adjusting eye velocity to match target motion. Processed signals are then relayed to brainstem premotor neurons, which activate the ocular motor nuclei to generate smooth and coordinated pursuit eye movements (Krauzlis, 2004; Leigh & Zee, 2015).

3.4 NEUROPHYSIOLOGY OF VERGENCE

Characteristics of Vergence: Vergence movements are disconjugate eye movements that adjust the angle between the eyes to maintain binocular fixation on objects at varying distances. During convergence, both eyes move inward to focus on near objects, whereas during divergence, they move outward to focus on distant objects. These movements are essential for binocular vision, stereopsis, and accurate depth perception (Leigh & Zee, 2015; Kandel et al., 2021).

Neural Pathways: Vergence is controlled by a network involving the visual cortex, frontal eye fields, and mesencephalic reticular formation, which process retinal disparity and generate vergence commands. Premotor neurons in the supraoculomotor area project to the oculomotor nucleus to coordinate eye movements. The posterior cerebellar vermis and fastigial nucleus refine vergence accuracy and facilitate its coordination with saccadic eye movements. Vergence is closely linked with accommodation and pupillary constriction as part of the near triad reflex (Gamlin, 2002; Leigh & Zee, 2015).

Integration of Saccades, Pursuits, and Vergence: Although saccades, smooth pursuits, and vergence are distinct eye movement systems, they share common cortical, brainstem, and cerebellar pathways. Their coordinated activity allows smooth transitions between gaze shifts and continuous tracking of objects at different distances, ensuring stable binocular vision during everyday visual tasks (Leigh & Zee, 2015; Robinson, 1981).

Clinical Manifestations of Dysfunction: Disorders affecting ocular motor pathways produce characteristic abnormalities. Lesions of the frontal eye fields or superior colliculus impair saccade initiation, while brainstem lesions involving the paramedian pontine reticular formation (PPRF) or rostral interstitial nucleus of the medial longitudinal fasciculus (riMLF) cause horizontal or vertical gaze palsies. Cerebellar disease results in saccadic dysmetria and impaired smooth pursuit, whereas vergence disorders, such as convergence insufficiency and internuclear ophthalmoplegia, commonly present with diplopia, reading difficulties, and reduced depth perception (Leigh & Zee, 2015).

Diagnostic Approaches: Evaluation of ocular motor function includes assessment of saccadic accuracy,

smooth pursuit, and the near point of convergence. Objective investigations such as infrared eye tracking, electrooculography (EOG), and functional MRI (fMRI) provide quantitative analysis of eye movements and help differentiate central from peripheral ocular motor disorders (Carpenter, 1988; Kandel et al., 2021).

Rehabilitation and Therapeutic Strategies: Management focuses on restoring ocular motor function through prism correction, orthoptic exercises, and vision therapy to improve vergence, pursuit, and binocular coordination. Neuroplasticity-based rehabilitation, including repetitive eye movement training, perceptual learning, and virtual reality-based therapy, has shown promising results in enhancing ocular motor performance and functional visual outcomes (Ciuffreda et al., 2009; Leigh & Zee, 2015).

3.5 FUTURE DIRECTIONS

Future research is focused on enhancing ocular motor rehabilitation through non-invasive brain stimulation, including transcranial magnetic stimulation (TMS) and transcranial direct current stimulation (tDCS), to improve neural control of eye movements. Advances in artificial intelligence (AI)-based eye tracking, virtual reality, and personalized rehabilitation programs are expected to provide more effective and individualized treatment for ocular motor disorders. Further research into the interaction between cognitive processes and ocular motor control will improve the understanding and management of neuro-ophthalmic diseases (Kandel et al., 2021; Leigh & Zee, 2015).

3.6 CONCLUSION

The neurophysiology of saccades, smooth pursuits, and vergence demonstrates the coordinated interaction of sensory processing, motor control, and cerebellar adaptation required for stable binocular vision. These eye movements rely on integrated cortical, brainstem, and cerebellar networks to maintain accurate gaze and visual stability. Advances in neuroimaging, eye-tracking technology, and neurorehabilitation continue to improve the diagnosis and treatment of ocular motor disorders, enhancing visual function and patient outcomes.

3.7 BIBLIOGRAPHY

1. Leigh RJ, Zee DS. The neurology of eye movements. 5th ed. New York: Oxford University Press; 2015.
2. Carpenter RHS. Movements of the eyes. 2nd ed. London: Pion; 1988.
3. Kandel ER, Koester JD, Mack SH, Siegelbaum SA. Principles of neural science. 6th ed. New York: McGraw-Hill Education; 2021.
4. Wurtz RH, Goldberg ME. The neurobiology of saccadic eye movements. Amsterdam: Elsevier; 1989.
5. Scudder CA, Kaneko CRS, Fuchs AF. The brainstem burst generator for saccadic eye movements: a modern synthesis. Exp Brain Res. 2002;142(4):439–62.
6. Sparks DL. The brainstem control of saccadic eye movements. Nat Rev Neurosci. 2002;3(12):952–64.
7. Robinson DA. Control of eye movements. Annu Rev Neurosci. 1981;4:463–503.
8. Krauzlis RJ. Recasting the smooth pursuit eye movement system. J Neurophysiol. 2004;91(2):591–603.
9. Ilg UJ. The role of areas MT and MST in smooth pursuit eye movements. Cereb Cortex. 2008;18(6):1383–94.
10. Thier P, Ilg UJ. The neural basis of smooth pursuit eye movements. Curr Opin Neurobiol. 2005;15(6):645–52.
11. Gamlin PDR. Neural mechanisms for the control of vergence eye movements. Ann N Y Acad Sci. 2002;956:264–72.
12. Judge SJ, Cumming BG. Neurons in the monkey midbrain with activity related to vergence eye movement and accommodation. J Neurophysiol. 1986;55(5):915–30.
13. Mays LE. Neural control of vergence eye movements: convergence and divergence neurons in the midbrain. J Neurophysiol. 1984;51(5):1091–108.
14. Ramat S, Straumann D, Zee DS. Gaze holding and the neural integrator. J Vestib Res. 2008;18(4):207–19.
15. Ciuffreda KJ, Ludlam DP, Kapoor N. Clinical oculomotor training in traumatic brain injury. Optometry. 2009;80(8):379–90.