Chapter-7.

TRAUMATIC BRAIN INJURY AND POST-VISUAL SYNDROMES

Dr Partha Chowdhury, Dr Brinda Shah

ABSTRACT

Traumatic brain injury is a multifaceted neurological condition with prevalent consequences across cognitive, motor and sensory domains. Among these, visual dysfunction is one of the most predominant yet underappreciated sequalae. Post visual syndromes subsequent TBI manifest through a spectrum of disorders which includes oculo motor dysfunction, cortical visual impairment, visual field deficiency and visuo perceptual anomalies. These deficiencies arise from diffuse axonal injury, cortical and subcortical lesions, metabolic compromise and maladaptive neural reorganization. Since vision reinforces navigation, learning and functional independence, impairment leads to thoughtful disability and psychological burden. This chapter provides as in-depth assessment of the neurobiological keystones, clinical manifestations, diagnostic approaches and rehabilitation approaches for visual dysfunction after TBI, while prominently showing future directions in neuro optometric care.

Keywords: Traumatic brain injury; Visual dysfunction; Neuro-optometric rehabilitation; Cortical visual impairment.

7.1 INTRODUCTION

Traumatic brain injury (TBI) is recognized as a leading cause of acquired disability worldwide, primarily affecting young adults, military veterans, and the elderly (Greenwald et al., 2012). Advances in critical care management have increased survival rates, but they have also revealed the long-term burden of persistent neurological sequelae. Vision, as the predominant sensory modality, is highly vulnerable to disruption following trauma (Ventura et al., 2014). The intricate architecture of the visual system, spanning from the retina through subcortical pathways to higher cortical centers, means that even minor injuries can cause significant visual dysfunction (Mendoza & Foundas, 2007; Zasler et al., 2022). Post-TBI visual syndromes contribute to disability in both obvious and subtle ways. Individuals may present with hemianopia that directly compromises mobility, or they may suffer from oculomotor fatigue, visual motion hypersensitivity, or cortical visual impairment (CVI), which interrupt daily living and academic or occupational functioning (Kapoor & Ciuffreda, 2002; Rowe et al., 2019). Unfortunately, many of these disturbances remain underdiagnosed due to their variable presentation and overlap with post-concussive cognitive symptoms (Goodrich et al., 2014). Distinguishing and addressing post-TBI visual sequelae is, therefore, a critical component of comprehensive neurorehabilitation (Pappadis et al., 2017).

7.2 NEUROBIOLOGICAL BASIS

The visual consequences of TBI arise from a convergence of functional and structural disruptions within the brain. Diffuse axonal injury (DAI) is a hallmark feature, occurring when shear forces from acceleration-deceleration impact disrupt white matter tracts (Zasler et al., 2022). The optic radiations, corpus callosum, and long association fibers connecting parietal, occipital, and temporal lobes are particularly susceptible, leading to impaired integration of visual information (Ventura et al., 2014). Direct cortical damage to the occipital lobe results in visual field deficits, whereas parietal injury produces visuospatial neglect and temporal lobe trauma disturbs object recognition (Mendoza & Foundas, 2007). Subcortical structures including the lateral geniculate nucleus (LGN) and superior colliculus may also be affected, impairing reflexive

visual responses. Moreover, brainstem or midbrain trauma can involve ocular motor nuclei and supranuclear pathways, leading to gaze palsy, diplopia, and vergence dysfunction (Ciuffreda et al., 2007). Secondary metabolic and vascular changes like ischemia, hypoxia, and excitotoxicity further exacerbate neuronal loss within visual circuits (Sabel et al., 2011).

7.3 CLINICAL MANIFESTATIONS

Visual dysfunction following traumatic brain injury (TBI) presents with a broad spectrum of clinical manifestations that reflect the complex involvement of cortical, subcortical, and brainstem visual pathways. The type and severity of symptoms vary according to the location and extent of neural injury, ranging from subtle ocular motor abnormalities to profound cortical visual impairment. These deficits significantly interfere with reading, mobility, occupational performance, and activities of daily living, thereby reducing overall quality of life. One of the most common manifestations of TBI is oculomotor dysfunction, which affects the coordination and control of eye movements. Patients frequently exhibit impaired saccades, reduced smooth pursuit, convergence insufficiency, accommodative dysfunction, and fixation instability. These abnormalities result in poor tracking of moving objects, blurred near vision, difficulty reading, visual fatigue, headaches, and intermittent diplopia. Convergence insufficiency is particularly prevalent following mild traumatic brain injury and post-concussion syndrome, contributing substantially to persistent visual discomfort and reduced reading efficiency (Ciuffreda et al., 2007; Capó-Aponte et al., 2012; Hunfalvay et al., 2020). Visual field defects are another important consequence of traumatic brain injury and commonly result from damage to the optic radiations, occipital cortex, or posterior visual pathways. Depending on the location of injury, patients may develop homonymous hemianopia, quadrantanopia, or localized scotomas that compromise peripheral vision and spatial awareness. These deficits increase the risk of falls, impair driving performance, reduce environmental awareness, and negatively affect independent mobility.

Although some spontaneous improvement may occur through cortical adaptation, many individuals require compensatory visual scanning strategies to improve functional performance (Suchoff et al., 2008). Cortical visual impairment (CVI) represents a higher-order visual disorder in which visual processing is impaired despite structurally normal eyes and intact anterior visual pathways. The condition arises from injury to the occipital cortex or association visual areas and primarily affects the dorsal ("where") and ventral ("what") visual processing streams. Patients may experience difficulty recognizing faces or objects, interpreting complex visual scenes, maintaining visual attention, and integrating visual information with cognitive function. Consequently, activities such as reading, navigation, and object localization become significantly impaired despite preserved visual acuity (Zasler et al., 2022). Many individuals with TBI also develop photosensitivity and visual motion hypersensitivity, particularly following concussion or diffuse axonal injury. Increased sensitivity to bright light often produces headaches, dizziness, ocular discomfort, nausea, and reduced visual endurance. Similarly, visually complex or dynamic environments, including crowded public spaces, scrolling computer screens, or moving traffic, may provoke disorientation, imbalance, and motion intolerance. These symptoms frequently overlap with post-concussion syndrome and substantially limit social participation, occupational performance, and community reintegration (Alnawmasi et al., 2022; Capó-Aponte et al., 2012). In addition, visuospatial and visuoperceptual deficits commonly occur following injury to the parietal and occipital association cortices. Patients may demonstrate impaired depth perception, spatial neglect, reduced visual-motor integration, impaired hand-eye coordination, and difficulties with spatial orientation. These higher-order visual deficits interfere with balance, navigation, reaching movements, driving, and the execution of complex daily tasks, creating significant safety concerns and increasing dependence on caregivers (Greenwald et al., 2012).

7.4 DIAGNOSTIC APPROACHES

The evaluation of visual dysfunction following traumatic brain injury requires a comprehensive, multidisciplinary approach that integrates detailed clinical assessment, neuro-optometric examination, neuroimaging, electrophysiological testing, and cognitive visual evaluation. Accurate diagnosis is essential for identifying the specific visual deficits, determining their underlying neuroanatomical basis, and developing individualized rehabilitation strategies. Assessment begins with a thorough clinical history focusing on symptoms such as blurred vision, diplopia, headaches, photosensitivity, reading difficulty, dizziness, visual fatigue, motion sensitivity, and problems with balance or navigation. The temporal relationship between symptom onset and traumatic injury, together with their impact on

functional activities, provides valuable information regarding the severity and progression of visual dysfunction (Richmond & Whittaker, 2020). Comprehensive neuro-optometric examination evaluates both sensory and motor components of vision. Testing includes measurement of visual acuity, refractive status, accommodation, convergence, binocular vision, ocular alignment, vergence ranges, fixation stability, saccadic eye movements, smooth pursuit, and vestibulo-ocular reflex function. These assessments help identify subtle ocular motor abnormalities that frequently remain undetected during routine ophthalmological examinations (Goodrich et al., 2014).

Objective investigations further enhance diagnostic accuracy. Automated perimetry is used to detect visual field defects, while contrast sensitivity testing identifies subtle impairments in visual perception that may not be reflected by standard visual acuity measurements. Electrophysiological investigations, particularly visual evoked potentials (VEP), assess the functional integrity of the visual pathways and assist in differentiating cortical from pre-cortical visual dysfunction. Advanced neuroimaging techniques provide valuable insights into the structural and functional consequences of traumatic brain injury. Structural magnetic resonance imaging (MRI) identifies focal cortical lesions, hemorrhage, and diffuse brain injury, whereas diffusion tensor imaging (DTI) evaluates white matter integrity and demonstrates diffuse axonal injury involving visual pathways. Functional magnetic resonance imaging (fMRI) further contributes by identifying altered cortical activation patterns and neural reorganization associated with visual recovery and compensatory neuroplasticity (Ventura et al., 2014). Evaluation of higher-order visual function is equally important, particularly in patients suspected of cortical visual impairment. Cognitive visual testing examines object recognition, facial recognition, visual attention, spatial perception, visual memory, executive visual processing, and visuospatial integration, thereby facilitating comprehensive assessment of cortical visual dysfunction and guiding appropriate rehabilitation strategies (Rowe et al., 2019).

7.5 REHABILITATION STRATEGIES

Rehabilitation of visual dysfunction following traumatic brain injury is individualized and multidisciplinary, combining neuro-optometric interventions, vision therapy, optical correction, compensatory training, and neuroplasticity-based rehabilitation to maximize functional recovery. Treatment aims to restore visual performance, improve ocular motor control, reduce symptoms, and enhance independence in daily activities. Vision therapy remains one of the primary rehabilitation approaches for post-traumatic visual dysfunction. Structured therapeutic exercises target impaired saccades, smooth pursuit, convergence, accommodation, fixation stability, and binocular coordination through repetitive, task-specific training designed to promote adaptive neuroplasticity. Regular therapy improves reading efficiency, reduces visual fatigue, enhances eye movement accuracy, and facilitates functional visual performance (Ciuffreda et al., 2007). Optical rehabilitation also plays a significant role in managing post-TBI visual disorders. Prism lenses are frequently prescribed to reduce diplopia, improve binocular alignment, compensate for ocular misalignment, and expand functional visual fields. Specialized tinted lenses, anti-glare coatings, and precision optical filters are beneficial for managing photosensitivity and improving visual comfort during prolonged visual activities.

Patients with visual field defects benefit from compensatory rehabilitation techniques that teach systematic visual scanning strategies to improve environmental awareness and mobility. In selected cases, computer-based visual field restitution therapy may stimulate residual cortical function and promote partial recovery of visual field awareness (Suchoff et al., 2008). Management of cortical visual impairment focuses on improving higher-order visual processing through multisensory integration training, cognitive rehabilitation, simplified environmental design, and task-specific functional exercises. These interventions encourage cortical reorganization while reducing visual complexity to facilitate more effective information processing (Zasler et al., 2022). Individuals experiencing motion hypersensitivity and vestibular symptoms often benefit from vestibulo-ocular rehabilitation, which combines gaze stabilization exercises, balance training, habituation therapy, and gradual exposure to visually stimulating environments. These interventions reduce dizziness, improve postural stability, and enhance tolerance to complex visual motion (Capó-Aponte et al., 2012).

Recent advances in neurorehabilitation have introduced innovative therapies aimed at enhancing cortical plasticity. Non-invasive brain stimulation techniques, including transcranial direct current stimulation (tDCS) and repetitive transcranial magnetic stimulation (rTMS), are being investigated as adjunctive therapies to facilitate neural recovery. Similarly, virtual reality (VR)-based rehabilitation provides immersive, interactive environments that

simulate real-world visual challenges while promoting adaptive neuroplasticity, improving visuomotor integration, and enhancing functional recovery (Sabel et al., 2011). Successful rehabilitation requires close collaboration among neuro-optometrists, neurologists, ophthalmologists, occupational therapists, physiotherapists, psychologists, and rehabilitation specialists to address the diverse visual, cognitive, and functional impairments associated with traumatic brain injury (Pappadis et al., 2017; Rowe et al., 2019).

7.6 FUTURE DIRECTIONS

Research into post-traumatic visual dysfunction is progressing rapidly with advances in neuroimaging, artificial intelligence, and neurorehabilitation technologies. Emerging neuroimaging biomarkers based on functional MRI, diffusion tensor imaging, and connectomic analysis are expected to improve early diagnosis, characterize cortical reorganization, and predict long-term visual outcomes following traumatic brain injury. Artificial intelligence and machine learning algorithms integrated with advanced eye-tracking systems have the potential to automate visual function assessment, identify subtle ocular motor abnormalities, and develop personalized rehabilitation programs tailored to individual patterns of neurological impairment. Virtual reality and augmented reality platforms continue to evolve as immersive rehabilitation tools that simulate realistic visual environments while providing real-time performance feedback and objective monitoring of recovery. Future rehabilitation is likely to adopt precision medicine approaches that combine neuroimaging, digital biomarkers, and individualized neuroplasticity-based interventions. Furthermore, greater integration of neuro-optometry within multidisciplinary neurorehabilitation teams will enhance comprehensive patient care and maximize functional recovery for individuals living with traumatic brain injury.

7.7 CONCLUSION

Traumatic brain injury frequently produces a diverse range of visual disorders that substantially impair functional independence, occupational performance, learning, mobility, and overall quality of life. Oculomotor dysfunction, visual field defects, cortical visual impairment, photosensitivity, motion intolerance, and visuospatial deficits reflect the vulnerability of the visual system to traumatic injury and the complex disruption of cortical and subcortical visual networks. Comprehensive clinical evaluation supported by advanced neuroimaging, neuro-optometric assessment, and electrophysiological testing is essential for accurate diagnosis and effective treatment planning. Individualized rehabilitation programs that integrate vision therapy, optical correction, compensatory strategies, neuroplasticity-based interventions, and emerging technologies such as virtual reality and non-invasive neuromodulation offer promising opportunities for functional recovery. Continued advances in neuroscience, neuroimaging, artificial intelligence, and multidisciplinary rehabilitation are expected to further improve the diagnosis, management, and long-term outcomes of visual dysfunction following traumatic brain injury, emphasizing the central role of vision in successful neurorehabilitation.

7.8 BIBLIOGRAPHY

1. Alnawmasi, M. M., Mann, D. L., & Lee, H.-S. (2022). The effect of mild traumatic brain injury on the visual processing of global form and motion. Brain Injury, 36(6), 743–752. https://doi.org/10.1080/02699052.2022.2063165
2. Capó-Aponte, J. E., Urosevich, T. G., Temme, L. A., Tarbett, A. K., & Sanghera, N. K. (2012). Visual dysfunctions and symptoms during the subacute stage of blast-induced mild traumatic brain injury. Military Medicine, 177(7), 804–813. https://doi.org/10.7205/milmed-d-12-00061
3. Ciuffreda, K. J., Kapoor, N., Rutner, D., Suchoff, I. B., Han, M. E., & Craig, S. (2007). Occurrence of oculomotor dysfunctions in acquired brain injury: A retrospective analysis. Optometry, 78(4), 155–161. https://doi.org/10.1016/j.optm.2006.11.011
4. Goodrich, G. L., Martinsen, G. L., Flyg, H. M., Kirby, J., Garvey, K. A., & Dawe, N. (2014). Development of a mild traumatic brain injury-specific vision screening protocol: A Delphi study. Journal of Rehabilitation Research and Development, 51(6), vi–xviii. https://doi.org/10.1682/JRRD.2013.10.0227
5. Greenwald, B. D., Kapoor, N., & Singh, A. D. (2012). Visual impairments in the first year after traumatic brain injury. Brain Injury, 26(11), 1338–1359. https://doi.org/10.3109/02699052.2012.706356
6. Hunfalvay, M., Roberts, C.-M., Murray, N., Tyagi, A., Barclay, K., & Bolte, T. (2020). Oculomotor behavior as a biomarker for differentiating Pediatric patients with mild traumatic brain injury and age matched controls. Frontiers in Behavioural Neuroscience, 14,

581819. https://doi.org/10.3389/fnbeh.2020.581819
7. Kapoor, N., & Ciuffreda, K. J. (2002). Vision disturbances following traumatic brain injury. Current Treatment Options in Neurology, 4(4), 271–280. https://doi.org/10.1007/s11940-002-0027-z
8. Mendoza, J. E., & Foundas, A. L. (2007). Clinical neuroanatomy: A neurobehavioral approach. Springer Science & Business Media.
9. Pappadis, M. R., Sander, A. M., Łukaszewska, B., Struchen, M. A., & Leung, P. (2017). The need for a holistic approach to post-traumatic brain injury visual impairment: A case report. Journal of Vocational Rehabilitation, 47(2), 235–246. https://doi.org/10.3233/JVR-170896
10. Richmond, P. W., & Whittaker, K. W. (2020). A practical approach to vision after brain injury. Practical Neurology, 20(3), 181–188. https://doi.org/10.1136/practneurol-2019-002384
11. Rowe, F. J., Hepworth, L. R., & Howard, C. (2019). Vision after acquired brain injury: A Delphi study of expert consensus on effective assessment and intervention. Disability and Rehabilitation, 41(21), 2521–2532. https://doi.org/10.1080/09638288.2018.1471168
12. Sabel, B. A., Henrich-Noack, P., Fedorov, A., & Gall, C. (2011). Vision restoration after brain injury and blindness: Therapeutic strategies and current achievements. Restorative Neurology and Neuroscience, 29(6), 395–411. https://doi.org/10.3233/RNN-2011-0613
13. Suchoff, I. B., Kapoor, N., Ciuffreda, K. J., Rutner, D., Han, E., & Craig, S. (2008). The frequency of occurrence, types, and characteristics of visual field defects in acquired brain injury: A retrospective analysis. Optometry, 79(5), 259–265. https://doi.org/10.1016/j.optm.2007.10.012
14. Ventura, R. E., Balcer, L. J., & Galetta, S. L. (2014). The neuro-ophthalmology of head trauma. The Lancet Neurology, 13(10), 1006–1016. https://doi.org/10.1016/S1474-4422(14)70111-5
15. Zasler, N. D., Katz, D. I., & Zafonte, R. D. (Eds.). (2022). Brain injury medicine: Principles and practice (3rd ed.). Demos Medical Publishing.